









Pharmaceutical Products



Antibiotics

Antibiotics are essential therapeutics in modern medicine. The two largest groups are Cephalosporins and Penicillins. In general, antibiotics are used in the treatment of bacterial infections.




AMOXICILLIN

500 mg powder for injection	10 x	
1 g powder for injection	50 x	
250 mg capsules	10 x 10	
500 mg capsules	1 x 10 10 x 10	
125 mg / 5 ml oral suspension (100 ml)	1 x	
250 mg / 5 ml oral suspension (100 ml)	1 x	

AMPICILLIN + SULBACTAM (TRO-SULPICIN)

1500 mg powder for injection/infusion	10 x	
---	------	---

AMPICILLIN

500 mg powder for injection/infusion	50 x	
1 g powder for injection/infusion	50 x	
500 mg capsules	10 x 10	

**BENZATHINE
PENICILLIN**

1.2 mega
powder for suspension for injection

50 x



2.4 mega
powder for suspension for injection

50 x



CEFAZOLIN

1 g
powder for injection/infusion

50 x



CEFOTAXIME

500 mg
powder and solvent for injection (water
for injection 10 ml)

1 + 1



1 g
powder and solvent for injection (water
for injection 10 ml)

1 + 1



1 g
powder and solvent for injection
(Lidocaine 1 %, 3.5 ml)

1 + 1



1 g
powder for injection/infusion

50 x



**CEFTAZIDIME
(TRO-CEFTADIME)**

1 g
powder for injection/infusion

10 x
50 x






2 g
powder for injection/infusion

10 x
50 x






CEFTRIAZONE (TRO-CEFTRIA)	1 g powder and solvent for injection (Lidocaine 1 %, 3.5 ml)	1 + 1	
	1 g powder and solvent for injection (water for injection 10 ml)	1 + 1	
	1 g powder for injection/infusion	50 x	
CEFUROXIME (TRO-XIME)	750 mg powder and solvent for injection (water for injection 10 ml)	1 + 1	
	750 mg powder for injection/infusion	50 x	
CHLORAMPHENICOL	1 % ophthalmic ointment (5 g)	1 x	
CIPROFLAXICIN (CIPRO-TRO)	2 mg / ml solution for infusion (100 ml)	1 x	
	500 mg film-coated tablets	1 x 10	
CLINDAMYCIN	300 mg / 2 ml injection	10 x	
	600 mg / 4 ml injection	5 x	


**CO-AMOXICLAV
(TRO-AMOXICLAV)**

600 mg powder for injection/infusion	10 x	
1200 mg powder for injection/infusion	10 x	
625 mg film-coated tablets	2 x 7 1 x 10	
1000 mg film-coated tablets	2 x 7 1 x 10	
156.25 mg / 5 ml oral suspension (70 ml)	1 x	
228.5 mg / 5 ml oral suspension (70 ml)	1 x	
312.5 mg / 5 ml oral suspension(70 ml)	1 x	
457 mg / 5 ml oral suspension(70 ml)	1 x	

CO-TRIMOXAZOLE

480 mg tablets	10 x 10	
960 mg tablets	10 x 10	
240 mg / 5 ml oral suspension (100 ml)	1 x	

**CRYSTALLINE
PENICILLIN**

1 mega powder for injection/infusion	50 x	
--	------	---

ERYTHROMYCIN STEARATE

250 mg
film-coated tablets

10 x 10



500 mg
film-coated tablets

10 x 10



125 mg / 5 ml
oral suspension (100 ml)

1 x



250 mg / 5 ml
oral suspension (100 ml)

1 x



FORTIFIED PROCAINE PENICILLIN

4 mega
powder for suspension for injection

50 x



GENTAMICIN

80 mg / 2 ml
injection

100 x



METRONIDAZOLE

5 mg / ml
solution for infusion (100 ml)

1 x



250 mg
tablets

10 x 10



500 mg
tablets

10 x 10



200 mg / 5 ml
oral suspension (100 ml)

1 x



STREPTOMYCIN SULPHATE

1 g
powder for injection

50 x



TETRACYCLINE HYDROCHLORIDE

1 %
ophthalmic ointment (5 g)

1 x



Analgesics/Antipyretics/Anti-inflammatories

Analgesics, antipyretics and anti-inflammatory drugs are pharmaceuticals for addressing diverse clinical conditions and are mainly used in the treatment of pain.

DICLOFENAC (TRO-DIFEN)

100 mg
suppositories

2 x 5
20 x 5



50 mg
suppositories

2 x 5
20 x 5



50 mg
tablets

100 x 10



75 mg / 3 ml
injection

10 x



DEXAMETHASONE SODIUM PHOSPHATE

4 mg/ml
injection

10 x



8 mg / 2 ml
injection

10 x



HYDROCORTISON SODIUM SUCCINATE

100 mg
powder and solvent for injection
(water for injection 2 ml)

1 +1



100 mg
powder for injection/infusion

50 x



IBUPROFEN

100 mg / 5 ml
oral suspension (100 ml)

1 x



400 mg
film-coated tablets

1 x 10
10 x 10



PARACETAMOL (TRO-PADOL)

500 mg / 5 ml
injection

5 x



10 mg / ml
solution for infusion (100 ml)

1 x



500 mg
tablets

10 x 10
100 x 10



125 mg
suppositories

2 x 5
20 x 5



250 mg
suppositories

2 x 5
20 x 5



500 mg
suppositories

2 x 5
20 x 5



120 mg / 5 ml
oral suspension (60 ml)

1 x



240 mg / 5 ml
oral suspension (100 ml)

1 x





Antidiabetics

Antidiabetics are used in the management of diabetes mellitus, a metabolic disorder marked by hyperglycemia. Depending on the severity, different types with different modes of action must be used for proper treatment.

METFORMIN (TRO-FORMIN)

500 mg
film-coated tablets

3 x 10



850 mg
film-coated tablets

3 x 10



1000 mg
film-coated tablets

3 x 10



Anti-ulcer Agents

Pharmaceutical products which are used for the treatment of peptic ulcer and gastroesophageal reflux disease (GORD).

OMEPRAZOLE

40 mg
powder and solvent for injection
(water for injection 10 ml)

1 + 1



40 mg
powder for injection/infusion

5 x



20 mg
capsules

7 x 2
10 x 10





Carrier Solutions

Carrier solutions are important in many treatments and diverse issues. They are often used in combination with other pharmaceutical products but can also have own therapeutic indications.

GLUCOSE

5 % intravenous infusion (500 ml)	1 x	
10 % intravenous infusion (500 ml)	1 x	

SODIUM CHLORIDE

0.9 % solution for infusion (500 ml)	1 x	
--	-----	--



Erectile dysfunction

Pharmaceutical products used for the treatment of erectile dysfunction support smooth muscle relaxation in blood vessels resulting in increased blood flow to the penis after sexual stimulation.

SILDENAFIL (TRO-VIAGILE)

100 mg film-coated tablets	1 x 4	
--------------------------------------	-------	--



Anticoagulants

Pharmaceutical products that are used to inhibit blood clotting.

ENOXAPARIN SODIUM

4000 IU (40 mg) / 0.4 ml

solution for injection in a
pre-filled syringe

2 x



The anticoagulant enoxaparin is a low molecular weight heparin that is used to inhibit blood clotting. Hence, the active substance is used in particular for the prevention and treatment of thromboses and embolisms. It is usually administered subcutaneously using a pre-filled syringe.

CARING FOR GLOBAL HEALTH

Visit our new website:

